Please check the examination details belo	ow before ente	ring your candidate information
Candidate surname		Other names
Centre Number Candidate		al GCSE
Monday 18 Novemb	er 202	24
Morning (Time: 1 hour 15 minutes)	Paper reference	4CH1/2C
Chemistry UNIT: 4CH1 PAPER: 2C		
You must have: Calculator		Total Marks

### **Instructions**

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- Fill in the boxes at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
  - there may be more space than you need.
- Show all the steps in any calculations and state the units.

### Information

- The total mark for this paper is 70.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each question.

## **Advice**

- Read each question carefully before you start to answer it.
- Write your answers neatly and in good English.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶





# The Periodic Table of the Elements

0	He helium	20 Neon 10	40 <b>Ar</b> argon 18	84 <b>Kr</b> krypton 36	131 <b>Xe</b> xenon 54	[222] <b>Rn</b> radon 86	t fully
7		19 <b>F</b> fluorine 9	35.5 CI chlorine 17	80 <b>Br</b> bromine 35	127 	[210] At astatine 85	orted but no
9		16 <b>O</b> oxygen 8	32 <b>S</b> sulfur 16	79 <b>Se</b> selenium 34	128 <b>Te</b> tellurium 52	[209] <b>Po</b> polonium 84	ive been rep
2		14 <b>N</b> nitrogen 7	31 <b>P</b> phosphorus 15	75 <b>As</b> arsenic 33	122 <b>Sb</b> antimony 51	209 <b>Bi</b> bismuth 83	rs 112-116 ha authenticated
4		12 <b>C</b> carbon 6	28 <b>Si</b> silicon 14	73 <b>Ge</b> germanium 32	119 <b>Sn</b> tin 50	207 <b>Pb</b> lead 82	Elements with atomic numbers 112-116 have been reported but not fully authenticated
ო		11 <b>B</b> boron 5	27 <b>AI</b> aluminium 13	70 <b>Ga</b> gallium 31	115 <b>In</b> indium 49	204 <b>T</b> thallium 81	nents with atc
				65 <b>Zn</b> zinc 30	112 <b>Cd</b> cadmium 48	201 <b>Hg</b> mercury 80	Elen
				63.5 <b>Cu</b> copper 29	108 <b>Ag</b> silver 47	197 <b>Au</b> gold 79	[272] <b>Rg</b> roentgenium 111
				59 nickel 28	106 <b>Pd</b> palladium 46	195 <b>Pt</b> platinum 78	[271]
				59 <b>Co</b> cobalt 27	103 <b>Rh</b> rhodium 45	192 <b>Ir</b> iridium 77	[268]
	1 <b>H</b> hydrogen			56 <b>Fe</b> iron 26	101 <b>Ru</b> ruthenium 44	190 <b>Os</b> osmium 76	[277] <b>Hs</b> hassium 108
				55 Mn manganese 25	[98] <b>Tc</b> technetium 43	186 <b>Re</b> rhenium 75	[264] <b>Bh</b> bohrium 107
		mass <b>ɔol</b> ıumber		52 <b>Cr</b> chromium 24	96 <b>Mo</b> molybdenum 42	184 <b>W</b> tungsten 74	[266] <b>Sg</b> seaborgium 106
	Key	relative atomic mass atomic symbol name atomic (proton) number		51 <b>V</b> vanadium 23	93 <b>Nb</b> niobium 41	181 <b>Ta</b> tantalum 73	[262] <b>Db</b> dubnium 105
		relativ <b>ato</b> atomic		48 <b>Ti</b> titanium 22	91 <b>Zr</b> zirconium 40	178 <b>Hf</b> hafnium 72	[261] <b>Rf</b> nutherfordium 104
				45 Sc scandium 21	89 <b>4</b> yttrium 39	139 <b>La*</b> lanthanum 57	[227] <b>Ac*</b> actinium 89
7		9 <b>Be</b> beryllium 4	24 <b>Mg</b> magnesium 12	40 <b>Ca</b> calcium 20	88 Sr strontium 38	137 <b>Ba</b> barium 56	[226] <b>Ra</b> radium 88
_		7 Li lithium 3	23 <b>Na</b> sodium 11	39 <b>×</b> potassium 19	85 <b>Rb</b> rubidium 37	133 <b>Cs</b> caesium 55	[223] <b>Fr</b> francium 87

<sup>\*</sup> The Lanthanides (atomic numbers 58-71) and the Actinides(atomic numbers 90-103) have been omitted.

Cu and Cl have not been rounded to the nearest whole number.



## **Answer ALL questions.**

Some questions must be answered with a cross in a box  $\boxtimes$ . If you change your mind about an answer, put a line through the box  $\boxtimes$  and then mark your new answer with a cross  $\boxtimes$ .

1 The table shows the composition of five species that are either atoms or ions.

Species	Number of protons	Number of neutrons	Number of electrons
А	1	0	1
В	3	3	3
С	3	4	3
D	9	10	10
E	13	14	13

(a) Use the table to answer these questions.

You may use each letter once, more than once, or not at all.

(i) Give the letter of the species that is a negative ion.

(1)

(ii) Give the letter of the species that has a full outer shell of electrons.

(1)

(iii) Give the letter of the species that has an atomic number equal to its mass number.

(1)




(b) Explain why species B and C are atoms of the same element with identical	
chemical properties.	(2)
(c) A proton has a mass of $1.6726 \times 10^{-24}$ g.	
A neutron has a mass of $1.6740 \times 10^{-24}$ g.	
Calculate the mass, in grams, of the nucleus of an atom of species C.	(2)
	\_/
mass =	<u>C</u>
(Total for Question 1 = 2	7 marks)



- 2 This question is about metals and metal compounds.
  - (a) All metals are malleable and good conductors of electricity.

The diagram shows the structure of a metal.



(i) Explain why metals are malleable.



(ii) Explain why metals are good conductors of electricity.


(2)



(k	<ul> <li>Two methods of obtaining a metal from a compound are electrolysis and extraction using carbon.</li> </ul>	
	Explain, without giving practical details, which method is most suitable for extracting iron from iron(III) oxide.	(2)
((	A student has a solution of iron(II) sulfate and a solution of iron(III) sulfate.	
	Describe a chemical test the student could use to identify which solution is iron(II) sulfate.	
		(2)
	(Total for Question 2 = 8 r	marks)

(1)

**3** A teacher uses this apparatus to separate a mixture of ethanol and water.



- (a) Give the name of this method of separation.
  - A chromatography
  - **B** crystallisation
  - **C** filtration
  - **D** fractional distillation



(b) Name the pieces of apparatus labelled X, Y and Z.	(3)
X	
Υ	
Z	
(c) Give a physical test that the teacher could do to find out if the ethanol produced is pure.	(2)
(Total for Question 3 = 6 ma	arks)

4	4 teacher	adds a	small piece	of lithium	to water an	d collects the	gas produced.
---	-----------	--------	-------------	------------	-------------	----------------	---------------

(a) (i) Give two observations when lithium is added to water.

(2)

2.....

(ii) The teacher adds a few drops of universal indicator to the solution at the end of the reaction.

Explain the final colour of the universal indicator.

(2)

(b) This is the equation for the reaction between lithium and water.

$$2Li(s) + 2H_2O(l) \rightarrow 2LiOH(aq) + H_2(g)$$

The teacher collects 550 cm<sup>3</sup> of hydrogen at rtp.

Calculate the mass, in grams, of lithium that the teacher added to the water.

[one mole of gas at rtp has a volume of 24000 cm<sup>3</sup>]

[for lithium,  $A_r = 7$ ]

[for carbon  $A_r = 12$  for hydrogen,  $A_r = 1$  for oxygen  $A_r = 16$ ]

(3)

mass = ..... g

(c) The teacher then adds a small piece of potassium to water.	
(i) Give one observation seen with potassium that is <b>not</b> seen with lithium.	(1)
(ii) Explain why potassium is more reactive than lithium.	
Refer to atomic structure in your answer.	(3)
(Total for Question 4 = 11	marks)



**5** A student does some titrations to find the volume of dilute nitric acid needed to exactly neutralise 25.0 cm<sup>3</sup> of sodium hydroxide solution.

This is the student's method.

- Step 1 add 25.0 cm<sup>3</sup> of sodium hydroxide solution to a conical flask
- Step 2 add three drops of methyl orange indicator
- Step 3 fill a burette with the acid
- Step 4 add acid from the burette until the indicator changes colour
- Step 5 record the volume of acid added
- (a) (i) Give the name of the apparatus that the student should use to measure the volume of sodium hydroxide solution in step 1.

(1)

(ii) Give the colour change seen in step 4.


(1)


(iii) Give a reason why the student does not use universal indicator in this titration.

(1)

(b) The student completes a rough titration and four accurate titrations.

The diagram shows the burette readings from the rough titration.





The table shows the student's results.

	Rough titration	Titration 1	Titration 2	Titration 3	Titration 4
Burette reading at end in cm <sup>3</sup>		21.80	22.85	21.75	24.10
Burette reading at start in cm <sup>3</sup>		0.50	0.15	0.25	0.10
Volume added in cm <sup>3</sup>		21.30	22.70	21.50	24.00

(i) Complete the table by adding the results from the rough titration.

Record the volumes to the nearest 0.05 cm<sup>3</sup>

(2)

(ii) Concordant results are results within 0.20 cm<sup>3</sup> of each other.

Use the concordant results from the table to calculate the mean volume of acid added.

(3)

mean volume of acid = ......cm<sup>3</sup>

(5)

(c) This is the equation for the reaction between dilute nitric acid and sodium hydroxide solution.

$$HNO_3(aq) + NaOH(aq) \rightarrow NaNO_3(aq) + H_2O(l)$$

After the titration, the student knows the volume of acid needed to neutralise 25.0 cm<sup>3</sup> of the sodium hydroxide solution.

Sodium nitrate decomposes at high temperatures.

Describe how the student could obtain pure, dry crystals of sodium nitrate from dilute nitric acid and sodium hydroxide solution.

(Total for Question 5 = 13 marks)

(4) 2	escribe how crude oil is separated into fractions by fractional distillation.	(4)
		(4)
	ome of the long-chain alkanes obtained from fractional distillation are cracked, roducing shorter-chain alkanes and ethene.	
р		(2)
р	roducing shorter-chain alkanes and ethene.	(2)
р	roducing shorter-chain alkanes and ethene.	(2)
р	roducing shorter-chain alkanes and ethene.	(2)
р	roducing shorter-chain alkanes and ethene.	(2)
р	roducing shorter-chain alkanes and ethene.	(2)
) ————————————————————————————————————	roducing shorter-chain alkanes and ethene.	
) ————————————————————————————————————	roducing shorter-chain alkanes and ethene.  ) Give the conditions necessary for cracking to occur.	(2)
) ————————————————————————————————————	roducing shorter-chain alkanes and ethene.  ) Give the conditions necessary for cracking to occur.	
) ————————————————————————————————————	roducing shorter-chain alkanes and ethene.  ) Give the conditions necessary for cracking to occur.	
) ————————————————————————————————————	roducing shorter-chain alkanes and ethene.  ) Give the conditions necessary for cracking to occur.	



# (iii) Ethene reacts with bromine.

The equation shows the displayed formulae of the reactants and product.

The table shows some bond energies.

Bond	Bond energy in kJ/mol
C=C	612
C—C	348
С—Н	414
Br—Br	193
C—Br	276

Show that the molar enthalpy change,  $\Delta H$ , for this reaction is about –100 kJ/mol.

(3)

$$\Delta H = \dots kJ/mol$$

							<b>(T</b>	otal for (	Questio	n 6 = 13	marks)
				,							(2)
	Refer	to bor	nd ener	gies in y	our ans	swer.					
(iv)	Expla	in why	the rea	action b	etween	ethene	and bro	omine is e	exotherr	nic.	
											reaction between ethene and bromine is exothermic.

- 7 This question is about alcohols and carboxylic acids.
  - (a) These are the structural formulae of alcohol A and carboxylic acid B.

**Alcohol A** 

CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>OH

**Carboxylic acid B** 

CH<sub>3</sub>(CH<sub>2</sub>)<sub>n</sub>COOH

(i) Name alcohol A.

(1)

(ii) Draw the displayed formula of alcohol A.

(1)

(iii) Carboxylic acid B has a chain of carbon atoms with no branches.

The number of CH<sub>2</sub> units is represented by the letter n.

Calculate the value of n.

[for carboxylic acid B,  $M_r = 242$ ]

(3)

n –

(iv) Alcohol A and carboxylic acid B react together to form an ester.

Give the other product of the reaction.

(1)



(b) These are the structural formulae of dicarboxylic acid C and diol D.

**Dicarboxylic acid C** HOOCCH<sub>2</sub>CH<sub>2</sub>COOH

**Diol D** HOCH<sub>2</sub>CH<sub>2</sub>OH

(i) This is the equation for the reaction between dicarboxylic acid C and sodium hydroxide solution.

25.0 cm<sup>3</sup> of 0.150 mol/dm<sup>3</sup> sodium hydroxide solution is completely neutralised by 17.5 cm<sup>3</sup> of a solution of dicarboxylic acid C.

Calculate the concentration, in mol/dm<sup>3</sup>, of the solution of dicarboxylic acid C.

Give your answer to three significant figures.

(4)

concentration = ...... mol/dm<sup>3</sup>



(ii) Dicarboxylic acid C and diol D react to form a polyester.

Draw the displayed formula of the repeat unit of this polyester.

(2)

(Total for Question 7 = 12 marks)

**TOTAL FOR PAPER = 70 MARKS** 





